

# Modelling a set of carbon-rich AGB stars at high-angular resolution

G. Rau<sup>1</sup>, J. Hron<sup>1</sup>, C. Paladini<sup>2</sup>, B. Aringer<sup>3</sup>, K. Eriksson<sup>4</sup>, P. Marigo<sup>3</sup>, W. Nowotny<sup>1</sup>, and R. Grellmann<sup>5</sup><sup>1</sup>University of Vienna, Department of Astrophysics, Türkenschanzstrasse 17, A-1180 Vienna<sup>2</sup>Institut d'Astronomie et d'Astrophysique, Université libre de Bruxelles, Boulevard du Triomphe CP 226, B-1050 Bruxelles, Belgium<sup>3</sup>Astronomical Observatory of Padova – INAF, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy<sup>4</sup>Department of Physics and Astronomy, Division of Astronomy and Space Physics, Uppsala University, Box 516, 75120 Uppsala, Sweden<sup>5</sup>Physikalisches Institut der Universität zu Köln, Zùlpicher Str. 77, 50397 Köln - Germany

## Abstract

We compared spectro-photometric and interferometric observations of six carbon-rich AGB stars with a grid of self-consistent model atmospheres. The targets are: R Lep, R Vol, Y Pav, AQ Sgr, U Hya and X TrA. Please refer to the publication Rau et al. (subm) for further details on those findings.

We used VLTI/MIDI interferometric observations at high-angular resolution, and spectro-photometric data. We compared the observations with self-consistent dynamic models atmospheres from the Uppsala group (Eriksson *et al.*, 2014; Mattsson *et al.*, 2010). We found partly similar results as in Rau *et al.* (2015). The models can reproduce SED data well at wavelengths longward of  $1\ \mu\text{m}$ , and the interferometric observations between  $8\ \mu\text{m}$  and  $10\ \mu\text{m}$ . Shortwards of  $1\ \mu\text{m}$  in the SED, and longwards of  $10\ \mu\text{m}$  in the visibilities, we note discrepancies which could be due to a combination of effects: data- and model-related.

The models which fit the best the Miras are significantly extended, with a significant shell-like structure. On the contrary, the models best fitting the non-Miras are more compact, showing lower average mass-loss rate.

We derived stellar parameters from the fits:  $T_{\text{eff}}$ ,  $L_{\text{models}}$ ,  $M$ ,  $C/O$ ,  $\dot{M}$ . The results are in good agreement with values from the literature, within the uncertainties.  $T_{\text{eff}}$  agrees with the temperature derived from the angular diameter  $\theta_{(V-K)}$  and the bolometric luminosity from the SED fitting,  $L_{\text{bol}}$ , except for AQ Sgr. We discuss in the text the likely reasons.

Finally, the Rosseland diameter  $\theta_{\text{Ross}}$  and  $\theta_{(V-K)}$  (van Belle *et al.*, 2013) agree with each other better for the Miras targets than for the non-Miras, and we speculate that the reason for that resides in the episodic mass loss of the latter models.

We placed the stars in the H-R diagram (see Fig. 1), confronting them with evolutionary tracks. The main derived properties ( $L$ ,  $T_{\text{eff}}$ ,  $C/O$  ratios and stellar masses) from the model fitting results in good agreement with TP-AGB evolutionary calculations for carbon stars (Marigo *et al.*, 2013).

(project STARKEY, G.A. n. 615604). CP is supported by Belgian Fund for Scientific Research F.R.S.- FNRS. We thank the ESO Paranal team for supporting our VLTI/MIDI observations. This research has made use of the Jean-Marie Mariotti Center Aspro service <sup>1</sup> and of SIMBAD database, operated at CDS, Strasbourg, France, and NASA/IPAC Infrared Science Archive.

## References

- Eriksson, K., Nowotny, W., Höfner, S., Aringer, B., & Wachter, A. 2014, *A&A*, 566, A95.  
Marigo, P., Bressan, A., Nanni, A., Girardi, L., & Pumo, M. L. 2013, *MNRAS*, 434, 488.  
Mattsson, L., Wahlin, R., & Höfner, S. 2010, *A&A*, 509, A14.  
Rau, G., Paladini, C., Hron, J., Aringer, B., Groenewegen, M. A. T., *et al.* 2015, *A&A*, 583, A106.  
van Belle, G. T., Paladini, C., Aringer, B., Hron, J., & Ciardi, D. 2013, *ApJ*, 775, 45.

## Acknowledgments

This work is supported by the “Abschlussstipendium fellowship” of the University of Vienna, which GR thanks, and by the FWF project P23006. P.M. and B.A. acknowledge the support from the ERC Consolidator Grant funding scheme

<sup>1</sup>Available at <http://www.jmmc.fr/aspro>




Figure 1: AGB region of the H-R diagram (see also Rau et al., subm). The lines show evolutionary tracks with solar metallicity from Marigo *et al.* (2013). The numbers denote the mass values at the beginning of the thermal pulsing (TP)-AGB. In a dotted line, for better visibility, is displayed the track with  $2 M_{\odot}$ . Different symbols refer to temperatures and luminosities estimated via observations and through finding models best fitting with spectro-photometric-interferometric-observations. Several colors indicate the different targets. A typical error bar is shown in black in the lower left of the figure.